

The Effect of Triclosan on *In vitro* Embryonic Development in Rat ^{[1][2]}

Dicle ÇAYAN ^{1,a} Erdoğan UNUR ^{2,b} Mehtap NİSARİ ^{2,c} Dilara PATAT ^{3,d} Ertuğrul DAĞLI ^{4,e} Hilal AKALIN ^{5,f}

^[1] The project was supported by Erciyes University Scientific Research Projects Commission (TDK-2017-7083)

^[2] This study was presented "Effects of Triclosan on *in vitro* Embryonic Rat Development" as oral presentation in 1st International Mediterranean Anatomy Congress, 6-9 September 2018, Konya, Turkey

¹ Niğde Ömer Halisdemir University, Niğde Zübeyde Hanım School of Health, Department of Nursing Management, TR-51100 Niğde - TURKEY

² Erciyes University, Faculty of Medicine, Department of Anatomy, TR-38280 Kayseri - TURKEY

³ Lokman Hekim University, Faculty of Medicine, Department of Anatomy, TR-06510 Ankara - TURKEY

⁴ Ahi Evran University, Faculty of Medicine, Research Hospital, TR-40200 Kırşehir - TURKEY

⁵ Erciyes University, Faculty of Medicine, Department of Medical Genetics, TR- 38280 Kayseri - TURKEY

ORCID: ^a 0000-0001-7233-063X; ^b 0000-0003-2033-4350; ^c 0000-0002-1126-7478; ^d 0000-0001-5237-4846; ^e 0000-0002-5331-1289

^f 0000-0002-2580-836X

Article ID: KVFD-2020-23873 Received: 07.01.2020 Accepted: 23.07.2020 Published Online: 24.07.2020

How to Cite This Article

Çayan D, Unur E, Nisari M, Patat D, Dağlı E, Akalın H: The effect of triclosan on *in vitro* embryonic development in rat. *Kafkas Univ Vet Fak Derg*, 26 (5): 595-602, 2020. DOI: 10.9775/kvfd.2020.23873

Abstract

Triclosan (TCS) is a broad spectrum antimicrobial agent showing its effect by deactivating the fatty acid synthesis of bacteria. The aim of this study was to investigate the effects of TCS on *in vitro* embryonic development in rats and to determine the levels of caspases 2, 7, 8, and 9 inducing cells to apoptosis through gene expression. According to the TCS dose added to the culture whole rat serum, 3 experimental groups and a control group were formed with each including 10 embryos. After 48 h culturing period, embryos were subjected to morphological scoring for developmental evaluation. The levels of caspases 2, 7, 8, and 9 were measured by performing gene expression on 40 embryos. Significant decreases were obtained in all parameters of morphological scoring in the experimental groups as the dose of TCS increased. While the caspase-2 level showed a significant decrease among the groups and was inversely proportional to the level of TCS, the caspase-9 level showed a significant increase among the groups and was directly proportional to the level of TCS. In conclusion, TCS was determined to cause apoptosis in the cells via the intrinsic pathway during pregnancy period and lead to embryonic growth retardation, which increased with increased dose of TCS.

Keywords: *Triclosan, Rat embryo culture, Apoptosis, Gene expression, Caspase*

Triklosanın *In vitro* Embriyonik Rat Gelişimi Üzerine Etkisi

Öz

Triklosan, bakterilerin yağ asidi sentezini bozarak etkisini gösteren geniş spektrumlu bir antimikrobiyal ajandır. Bu çalışmanın amacı, triklosanın ratlarda *in vitro* embriyonik gelişim üzerindeki etkisini araştırmak ve gen ekspresyon yöntemi ile hücreleri apoptoza sürükleyen kaspaz 2, 7, 8 ve 9 değerlerini belirlemektir. Kültür şişesine eklenen triklosan dozuna göre, kontrol ve 3 deney grubu oluşturuldu. 48 saatlik kültür periyodundan sonra embriyolar gelişimsel değerlendirme amacıyla morfolojik skorlamaya tabi tutuldu. Ardından 40 embriyoya gen ekspresyonu yapılarak kaspaz-2, kaspaz-7, kaspaz-8 ve kaspaz-9 değerleri ölçüldü. Morfolojik skorlamaya ait tüm parametrelerde istatistiksel olarak anlamlı bir gerileme tespit edildi. Kaspaz-2 değerlerinin triklosan miktarı ile ters orantılı bir şekilde, gruplar arasında istatistiksel olarak anlamlı bir azalma gösterdiği, kaspaz-9 değerlerinin ise triklosan miktarıyla orantılı bir biçimde istatistiksel olarak anlamlı bir artış gösterdiği tespit edildi. Bu verilere göre, gebelik döneminde triklosan kullanımının, hücrelerde intrinsik yol aracılığıyla apoptoza yol açarak, embriyonik gelişme geriliğine neden olduğu, bu durumun doza bağlı olarak arttığı belirlendi.

Anhtar sözcükler: *Triklosan, Rat embriyo kültürü, Apoptoz, Gen ekspresyonu, Kaspaz*

INTRODUCTON

Triclosan (TCS) is a fat-soluble, broad-spectrum antimicrobial agent produced in laboratory in 1966. The area

of its usage was first expanded into antibacterial soaps and then into products such as toothpastes, cosmetics, and deodorants in the following years ^[1]. TCS can pass the biological barriers because of its lipophilic property



Correspondence



+90 533 4507645 Fax: +90 388 2112813



dcayan12@gmail.com

and can aggregate in living organisms [2]. The target organs have been reported to be the liver, spleen, brain, heart, reproductive system, and the immune system. Compared with other well-identified target organs, high concentrations of TCS up to 7 ng/mL can be detected in the placenta, cord blood, and amniotic fluid, which directly indicate a high risk of adverse effects on the embryo in an utero [3]. TCS is a relatively small compound with a molecular weight of 289.5 g/mole [4]. Owing to its molecular weight, TCS can pass the placenta and affect the embryo. TCS metabolism mainly occurs in the liver; however, it can also be metabolized in the skin in small amounts [5]. TCS is metabolized to glucuronide and sulfate conjugates (phase II metabolism) and primarily excreted by urine [6]. As a result of metabolism, it passes to body fluids, breast milk, and urine [7,8] and then merges into sewerage system via wastes, which causes wide contamination in the environment, such as in drinking water and surface waters [1,9]. Therefore, people using TCS-containing products are directly at risk and individuals drinking water contaminated with TCS are indirectly at risk. Many cell culture studies have shown that TCS is cytotoxic to human hepatocytes and cardiotoxic to cardiomyocytes, has neurodegenerative effects on rat neural stem cells, and shows strong embryotoxic effects on bone development [3,10-12]. However, only a few studies have reported its embryotoxic effects on embryo culture [4,13,14]. Moreover, the exact mechanism of the effects of TCS on early embryonic development is still not well understood. In a study with zebrafish embryos, growth retardation has been detected in the embryos exposed to 300 µg/L of TCS [13]. Another study has also demonstrated embryotoxic effects of TCS at high doses (250 µg/L) in zebrafish embryos, such as abnormal phenotypes, short tails, heart edema, and decreased hatching rate [4]. Developmental parameters are proved to be excellent indicators of TCS toxicity [4].

Developmental delay in babies has made the reliability of drugs and chemicals used in pregnancy period a current issue. The European Center for the Validation of Alternative Methods, established in 1991, has accepted the Rat Whole-Embryo Culture Test as one of the methods used to determine the teratogenic and toxic effects of chemical substances [15]. Therefore, this test is a preferred method in toxicology studies [16-19]. Previously, it was suggested that toxic substances triggered pregnancy loss, embryonic death or structural abnormalities by affecting the mechanisms that regulated normal embryonic development. However, today, it is known that apoptosis plays an important role in embryogenesis with cell proliferation and cell differentiation and that toxic substances disrupt the apoptosis pathway in the embryonic organs and cause malformations [20]. Apoptotic stimulus leads to intracellular activation of caspases. Pathways that activate initiator caspases vary with an apoptotic stimulus and there are two different pathways; extrinsic and intrinsic pathways. Extrinsic pathway is driven by caspase-8 and

caspase-7. Intrinsic pathway, which is also called classical or mitochondrial pathway, is the major route to apoptotic death in mammalian cells. This step triggers caspase-9 and caspase-2. These caspases are not specifically present in the tissue [21,22].

In the present study, we aimed to investigate the direct toxic effects of TCS on embryonic growth and development in cultured rat embryos and to evaluate its possible genotoxic effects by determining the potential role of apoptosis on the toxic effects of TCS using the gene expression method. Real-time PCR is the most sensitive molecular method with high sensitivity (10^5 - 10^6), which has the risk of contamination and has a short lead time. The activations of caspase-2, caspase-7, caspase-8, and caspase-9, which may adversely affect cell survival, were measured to explore the mechanisms underlying the effects of TCS in embryo.

MATERIAL and METHODS

The study was approved by the Animal Care and Use Committee (Ethics Committee) of Erciyes University (date: 13.01.2016 and No: 16/012). All procedures throughout the study were carried out in accordance with the ethical issues. Wistar rats were obtained from the Clinical and Experimental Research Center in the Medical Faculty of Erciyes University.

Chemicals

TCS used in the study was provided from Sigma-Aldrich (CAS No 3380-34-5) (Lot# LRAA9502) and GAPDH (Lot# 0000054846), caspase-2 (Lot# 0000054850), caspase-7 (Lot# 0000054845), caspase-8 (Lot# 0000054847), and caspase-9 (Lot# 90017681) were obtained from Roche.

Embryo Culture and Morphological Scoring

Female rats approximately 4-10 months of age and weighing 150-250 g were paired with their male partners in cages at about 5.00 pm and left overnight. On the next day morning around 8 am, the female rats were examined regarding the presence of vaginal plugs as an indication of mating and thereby fertilization. The female rats with vaginal plug were considered as 0.5 day pregnant at noon. The conceptuses were dissected from the uteri, decidua and Reichert's membranes by general anesthesia on day 9.5 of gestation after the blood samples of the females were collected from the abdominal aorta. The conceptuses were explanted into whole embryo culture by the method of New [23].

In order to assess the toxic effect of TCS on embryonic growth, the embryos were divided into four groups (each consisting 10 embryos); three experimental groups and one control group. In the literature, no study of TCS dose to be applied in embryo culture was found. Therefore, the

administration dosages of TCS were determined according to the data gained from previous studies and the reference dose was accepted as 300 ng/mL [4,13,24]. The control group embryos were cultured in whole rat serum (WRS) and the experimental groups were cultured in WRS containing 100 ng/mL, 200 ng/mL, and 300 ng/mL of TCS, in accordance with the technique developed by New [23]. Using this method, the effects of TCS on *in vitro* embryonic development during the early organogenesis period (between 9.5 and 11.5 days) were evaluated. Moreover, the embryonic development was also compared between the embryos of control and experimental groups morphologically on day 11.5 (after 48 h of culturing period) via morphologic scoring system [14]. In this scoring system, taking the growth and differentiation of different embryological features into consideration, 11.5 day embryos were evaluated by 17 parameters; each parameter was divided into 6 stages; and each stage was scored with a numerical value between 0 and 5. As morphological parameters, mean yolk sac diameter; yolk sac vessel development; allantois development; embryonic flexion; heart and caudal neural tube development; hind-brain, midbrain and forebrain development; development of eye, ear, nose and pharynx; maxillary and mandibular processes; differentiation fore and hind limbs; crown-rump length; and somite number were evaluated.

Gene Expression Stages

Tissue specific RNA isolation was not possible in immature embryos due to TCS toxicity. Therefore, we used the whole embryo to perform RNA isolation of caspases that are not specific to any tissue.

RNA isolation and cDNA synthesis: Rat embryos were placed into 500 μ L of TriPure Isolation Reagent (Roche Applied Science, Basel, Switzerland) for RNA isolation. Total RNA isolation was performed using the protocol for High Pure RNA Tissue Isolation Kit (Roche Applied Science, Mannheim, Germany). Qualification and quantification of RNA samples was performed using the Nanodrop 1000 Spectrophotometer (Thermo Fisher Scientific, Dreieich, Germany). RNA concentrations were assessed by optical density measurement at 260 nm and 280 nm and purity was determined by the ratio of 260/280 nm. Synthesis of cDNA from the total RNA (100 ng) was performed using random hexamer as a primer via the Transcriptor High Fidelity cDNA Synthesis Kit (Roche Applied Science, Mannheim, Germany) according to the manufacturer's protocol.

Quantitative real-time polymerase chain reaction (PCR) analysis:

From the obtained cDNA samples, mRNA expression levels of caspase-2, caspase-7, caspase-8, and caspase-9 genes, which have roles in apoptotic pathways, were investigated using quantifier real-time PCR via LightCycler® 480 II device (Roche Diagnostics, Mannheim, Germany). The primer sequences specific to the cDNAs of the investigated genes and the Universal Probe Library (UPL) probe numbers are presented in Table 1. Amplifications were performed in the total reaction volume of 20 μ L using cDNA, mRNA-specific primers, UPL probe, LightCycler® probe master mixture (Roche Diagnostics, Mannheim, Germany), and distilled water according to the PCR cycling program: 95°C for 10 min, then 45 cycles of 95°C for 10 sec, 60°C for 30 sec, followed by 72°C for 10 sec. The level of mRNA expression of glyceraldehyde-3-phosphate dehydrogenase (GAPDH) was accepted as reference for normalizing the amounts of caspase-2, caspase-7, caspase-8, and caspase-9 gene expression. The procedure was repeated three times for each concentration of TCS. The expression levels of target genes were calculated using the relative quantitation method using the software program of the LightCycler® 480 II device (Table 1).

Statistical Analysis

Data were analyzed using the IBM SPSS Statistics for Windows, Version 22.0 (IBM Corp., Armonk, NY, USA). The conformity of variables to normal distribution was assessed using the Shapiro-Wilk test, Q-Q plot, and histogram graphs. In multiple independent group comparisons, one-way analysis of variance was used for normally distributed numerical variables. In post-hoc comparisons, Tukey's test was used for normally distributed variables. The Pearson's correlation analysis was performed to determine the correlation between the levels of caspases and the morphological parameters. A P value <0.05 was considered statistically significant.

RESULTS

There was severe growth retardation in the embryos of the experimental groups as compared to the control group (Fig. 1). According to the morphological scoring system, the retardation of embryonic growth and development was increased as the dose of TCS increased (Fig. 2). The lower morphological scores were related with the poor

Table 1. Assay ID numbers, Universal probe number and primer sequences of genes used in expression study

Gene Symbol	Assay ID	Forward Primer Sequence	Reverse Primer Sequence	UPL No
Casp2	506708	5'-GGAAATGAGGGAGCTAATCCA-3'	5'-GGCAGCAGGTTGAGGAGTT-3'	129
Casp7	500441	5'-CTCTTGCGCAAAGATGCAG-3'	5'-AGCAGGCTGAGTTGCTGTG-3'	110
Casp8	504044	5'-GCCTGAGGGAAAGATGTCT-3'	5'-TCACATCATAGTTCACGCCAGT-3'	89
Casp9	502880	5'-CGACATGATCGAGGATATTCA-3'	5'-TGCCTCCCTCGAGTCTCA-3'	27
Gapdh	503799	5'-CATCGTGAAGGGCTCAT-3'	5'-CGCCACAGCTTCCAGAG-3'	158




Fig 1. Normally developing embryo cultured in WRS

yolk sac vessel development, failure of fusion of the neural folds; incomplete embryonic flexion; retardation in the development of otic, optic, and olfactory systems; branchial bars; maxillary and mandibular processes; and limbs (Table 2). The mean morphological scores of the embryos in the control groups was 62.80 ± 6.42 whereas those of the embryos cultured in the WRS containing 100 ng/mL, 200 ng/mL and 300 ng/mL of TCS were 30.4 ± 6.34 , 14.9 ± 6.55 , and 1.8 ± 3.01 , respectively. The difference in the morphological scores was statistically significant between




Fig 2. Embryos after 48 h culture. A: control group, B: 100 ng/mL triclosan group, C: 200 ng/mL triclosan group, D: 300 ng/mL triclosan group; Bar: 500 μ m

Table 2. Data variance analysis on morphological scoring between the control and experimental groups

Parameters	Groups				P
	Control	100 ng/mL Triclosan	200 ng/mL Triclosan	300 ng/mL Triclosan	
Yolk Sac Vessel Development	4.8 ± 0.422^a	3.3 ± 0.422^b	1.6 ± 0.823^c	0.3 ± 0.483^d	≤ 0.001
Allantois	3 ^a	1.9 ± 0.568^b	1 ± 0.471^c	0.3 ± 0.483^d	≤ 0.001
Flexion	4.9 ± 0.738^a	2.7 ± 0.949^b	1.5 ± 0.707^c	0.3 ± 0.483^d	≤ 0.001
Heart	4.9 ± 0.316^a	3 ± 0.471^b	1.7 ± 0.823^c	0.3 ± 0.483^d	≤ 0.001
Caudal Neural Tube	4.7 ± 0.483^a	2.4 ± 0.516^b	1.1 ± 0.568^c	0 ^d	≤ 0.001
Hindbrain	4.4 ± 0.516^a	2 ± 0.471^b	1.2 ± 0.632^c	0.2 ± 0.422^d	≤ 0.001
Midbrain	4.1 ± 0.738^a	1.8 ± 0.632^b	0.9 ± 0.316^c	0.2 ± 0.422^d	≤ 0.001
Forebrain	4.1 ± 0.738^a	1.6 ± 0.516^b	0.9 ± 0.316^c	0.2 ± 0.422^d	≤ 0.001
Otic System	4.1 ± 0.568^a	2.2 ± 0.789^b	1 ± 0.667^c	0 ^d	≤ 0.001
Optic System	4.5 ± 0.527^a	2.3 ± 0.675^b	1 ± 0.816^c	0 ^d	≤ 0.001
Olfactory System	2.7 ± 0.483^a	1.1 ± 0.316^b	0.4 ± 0.516^c	0 ^c	≤ 0.001
Pharyngeal Arch	2.9 ± 0.568^a	1.3 ± 0.483^b	0.6 ± 0.516^c	0 ^d	≤ 0.001
Maxillary Processes	2.5 ± 0.527^a	1.1 ± 0.316^b	0.5 ± 0.527^c	0 ^d	≤ 0.001
Mandibular Processes	2.5 ± 0.527^a	0.8 ± 0.422^{bc}	0.6 ± 0.516^{bc}	0 ^d	≤ 0.001
Forelimbs	2 ± 0.471	0.3 ± 0.483	0	0	≤ 0.001
Hindlimbs	1.9 ± 0.316	0.1 ± 0.316	0	0	≤ 0.001
Somits	4.8 ± 0.422^a	2.5 ± 0.707^b	0.9 ± 0.738^c	0 ^d	≤ 0.001
Total Morphological Score	62.8 ± 6.426^a	30.4 ± 6.346^b	14.9 ± 6.557^c	1.8 ± 3.011^d	≤ 0.001
Yolk Sac Diameter	3.590 ± 0.2132^a	2.68 ± 0.3011^b	1.61 ± 0.3872^c	1.15 ± 0.1434^d	≤ 0.001
Crown-Rump Length	2.95 ± 0.2173	1.97 ± 0.4057	0.86 ± 0.2459	-	≤ 0.001
Somit Number	26.8 ± 1.751	15.9 ± 4.067	7.1 ± 2.025	-	≤ 0.001

The data are as average and standard deviation. The same letters in the same row show similarity among groups, and different letters indicate differences among groups




Fig 3. Yolk sac vessel development. A: Control group, B: 100 ng/mL triclosan group, C: 200 ng/mL triclosan group, D: 300 ng/mL triclosan group; Bar: 1 mm




Fig 4. Effect of triclosan on total morphological score, yolk sac diameter mean, crown-rump length and somit number mean. Values are given as Mean and Standard Error, * P < 0.001

the control and experimental group ($P < 0.001$). In addition to the total morphological scores, yolk sac diameter, somit numbers, and crown-rump length were significantly lower in the experimental groups than those in the control group ($P < 0.001$ for each). While the mean yolk sac diameter was

3.59 ± 0.21 mm in the control group, it was 2.68 ± 0.30 mm, 1.61 ± 0.38 mm, and 1.15 ± 0.14 mm in the experimental groups exposed to 100 ng/mL, 200 ng/mL and 300 ng/mL of TCS, respectively (Fig. 3). The mean crown-rump length was 2.95 ± 0.2 mm in the control group; however, in the experimental groups, it gradually diminished as the dose of TCS increased (1.97 ± 0.40 mm, 0.86 ± 0.24 mm, and not measured for 100 ng/mL, 200 ng/mL and 300 ng/mL, respectively). The mean somit number was 26.8 ± 1.75 in the control group and it was also gradually diminished in the experimental groups as the dose of TCS increased (15.9 ± 4.06 , 7.1 ± 2.02 and, not measured for 100 ng/mL, 200 ng/mL and 300 ng/mL, respectively) (Fig. 4). The heart development values of embryos was also lower in the experimental groups (3 ± 0.47 , 1.7 ± 0.82 , and 0.3 ± 0.48 for 100 ng/mL, 200 ng/mL and 300 ng/mL, respectively) compared with that in the control group (4.9 ± 0.31) ($P < 0.001$) (Fig. 5).

The molecular mechanisms underlying the toxic effects of TCS on early development of rat (whole embryo culture) was assessed via quantitative real-time-PCR and gene expression. TCS significantly altered four caspase genes (mRNA expression levels) associated with apoptosis. When caspase levels of the experimental groups were compared with the control group, TCS caused upregulation of expression of caspase-7, caspase-8, and caspase-9 and downregulation of expression of caspase-2 (Table 3). There were significant differences between the experimental and control groups regarding the gene levels of caspase-2 and caspase-9 ($P < 0.001$) (Fig. 6). However, no significant difference was found between the groups in terms of the gene levels of caspase-7 ($P = 0.566$) and caspase-8 ($P = 0.396$). According to the correlation analysis of the levels of caspase-2, caspase-7, caspase-8, and caspase-9 with all study parameters, significant positive correlations were obtained between caspase-2 and morphological score, somit number, and yolk sac diameter. Additionally, there were significant negative correlations between caspase-9 and all morphological parameters ($P < 0.01$).

DISCUSSION

Triclosan has been widely used in personal care products or medical devices, such as sutures, owing to its antibacterial




Fig 5. Heart development. **A:** control group, **B:** 100 ng/ml triclosan group, Arrow: Heart; Bar: 500 μ m




Fig 6. Caspase levels of groups

Table 3. Mean variance analysis of experimental groups and control group

Caspase	Groups				P
	Control	100 ng/mL Triclosan	200 ng/mL Triclosan	300 ng/mL Triclosan	
Caspase 2	0.733±0.194 ^a	0.570±0.167 ^a	0.564±0.166 ^a	0.303±0.14 ^b	<0.001
Caspase 7	1.047±0.444	1.112±0.319	1.389±0.512	1.31±0.977	>0.05
Caspase 8	0.852±0.209	0.870±0.166	1.701±0.802	1.888±3.36	>0.05
Caspase 9	1.041±0.285 ^a	1.333±0.179 ^{a,b}	1.805±0.643 ^b	1.937±0.740 ^b	<0.001

The data are as average and standard deviation. Different letters in the same row indicate differences between groups

properties. It is known that TCS has potential risks to reproduction and development. Although the adverse effects of TCS on different organs have been studied for decades, the mechanism of its toxicity is still not well understood, especially during the stages of embryo development [4,13]. Whole embryo culture method is a valuable model for assessing the effects of toxic agents on early embryogenesis. In this method, rat embryos are cultured *in vitro* after the gestational day 9.5-11.5, which is a critical period for organogenesis in rats and corresponds to 3-6 weeks after fertilization in human embryos. However, to the best of our knowledge, there are no studies in the literature verifying the toxic responses to TCS during embryo development in rat embryo culture. The present study highlighted the effects of TCS on *in vitro* embryonic development using cultured rat embryos and demonstrated the negative effect of TCS on cell viability leading to induction of apoptosis.

The *in vitro* embryotoxic effect of TCS using embryo/larvae

of zebrafish (*Danio rerio*) was previously reported by Oliveira et al. [4]. Embryos were exposed to different doses of TCS (500 ng/mL, 700 ng/mL, and 900 ng/mL) for 6 days and embryotoxic effects (including delay in hatching, abnormal eye and body pigmentation, spinal malformations, cardiac edema, and body size smaller than normal) were observed [4]. In another study, Chen et al. [13] found no significant increase in the mortality after TCS treatment; however, 300 μ g/L of TCS caused developmental retardation in zebrafish embryos at 24 h post fertilization. Ho et al. [25] examined the morphology of zebrafish larvae after TCS exposure and they observed that the zebrafish embryos exposed to 250 ng/mL of TCS developed normally, similar to the control groups. They concluded that low-dose TCS exposure did not affect general embryonic development [14]. In the study by Guo et al. [26], TCS was administrated *in ovo* to evaluate its toxic effects in the chicken embryos. Tarsus length (mm) was significantly shorter in the embryos exposed to 10 μ g/g TCS compared with that in the embryos administered corn oil [26]. In another study, it was reported that exposure

to 8-10 μM TCS led to a high rate of developmental delay at 24 h post fertilization in zebrafish embryos [27]. In the present study, for the first time, effects of TCS on cultured rat embryos were examined. The results showed that the total embryonic growth was normal in the embryos cultured in the WRS only as opposed to the embryos cultured in the WRS containing TCS; this figure decreased depending on the dose of TCS. Embryonic retardations were observed in total embryonic growth, especially in the yolk sac diameter and vascularization, crown-rump length, somite number, body flexion, and neural tube development. These parameters were significantly lower in the experimental groups compared with the control group.

Researches have reported that TCS exposure can induce apoptosis in various biological systems [28-30]. In the present study, cultured embryos were also evaluated regarding apoptosis to investigate the potential role of apoptosis on the embryotoxic effects of TCS via caspases using the gene expression method. Since caspase-2, caspase-7, caspase-8, and caspase-9, which were measured to assess apoptosis, were present in all tissues, we performed RNA isolation on the whole embryo. Several studies have showed that caspase-2 is activated by DNA damage in the nucleolus [31,32]. Dubey et al. [29] reported that TCS induced apoptosis in human skin keratinocytes and responsible for DNA damage. In contrast, our data showed that TCS reduced the activity of caspase-2; if TCS had caused DNA damage, an increase in the level of caspase-2 would have occurred. Therefore, we could say that TCS did not cause DNA damage within the cell and that the decrease in caspase-2 level was associated with growth retardation. On the other hand, there is no data in the literature concerning the caspase-7 levels of TCS. Lamkanfi and Kanneganti [33] claimed that caspase-7 was related to inflammation; the activation of inflammatory caspase-1 and incorporation with caspase-7 could cause inflammation in the cell by the formation of a structure called inflammasome. Brentnall et al. [34] reported that caspase-9 and caspase-7 had different roles in intrinsic apoptosis. While intrinsic apoptosis resulted in the activation of caspase-9, caspase-7 did not have a role in intrinsic apoptosis but enabled apoptotic cell detachment [34]. In the present study, as the dose of TCS increased, caspase-7 levels increased; however, no significant difference was observed between the groups. This finding suggested that TCS added to the embryo culture medium did not cause inflammation within the cell. In the *in vitro* study investigating the apoptotic effect of TCS on mouse neocortical neurons, Szychowski et al. [28] evaluated the levels of caspase-8 and caspase-9 and they found that the level of caspase-8 significantly increased as the exposure duration to TCS and the concentration of TCS increased. However, they observed the increase in caspase-9 level in the groups at high concentration of TCS. Accordingly, the researchers claimed that non-cytotoxic concentrations of TCS caused apoptosis through the extrinsic pathway; however, long-term exposure of TCS activated intrinsic

pathway via caspase-9 by releasing cytochrome-c from the mitochondria [28]. In the present study, caspase-7 and caspase-8 levels increased as the dose of TCS increased; however, the difference was not statistically significant between the groups. Caspase-9 has a key role in the early stage of mitochondrial apoptosis pathway by activating downstream caspases and initiating apoptosis [35]. Li et al. [36] reported that methyl-TCS was a dominant transformation product of TCS, exposure to which induced the increased expression of caspase-9 on cell culture. In the present study, caspase-9 levels were found to be increased among the groups in proportion to TCS. It is known that TCS causes cell apoptosis via caspase-9.

In the present study, TCS significantly decreased all growth and developmental parameters in a dose-dependent manner in the embryos of the experimental groups compared with the control embryos. This finding suggests that TCS can activate the intrinsic pathway in the cell by passing directly through the cell membrane and that the intrinsic pathway is functional in the embryonic period.

In conclusion, the results of the present study revealed that TCS was a toxic agent and led to growth retardation by causing apoptosis through the intrinsic pathway without resulting in inflammation and direct DNA damage in the cell. Moreover, the current results highlighted the potential role of TCS as a strong embryotoxic in embryonic rats. These results would be an important source for the effects of TCS on embryonic development. Future studies isolating embryonic tissues and measuring the caspase levels by gene expression would also make contribution to investigate the effects of TCS on embryonic development.

CONFLICT OF INTEREST

The authors declare that there are no conflicts of interest.

AUTHOR CONTRIBUTIONS

D. Çayan and E. Unur conceived the ideas of the study and writing manuscript; D. Çayan, M. Nisari, D. Patat and E. Dağlı performed data collection and analysis; H. Akalın performed gene expression stages.

REFERENCES

1. Glaser A: The Ubiquitous TCS. *Pest You J*, 24 (3): 12-17, 2004.
2. Szychowski KA, Wnuk A, Rzemieniec J, Kajta M, Leszczyńska T, Wójtowicz AK: Triclosan-evoked neurotoxicity involves NMDAR subunits with the specific role of glun2a in caspase-3-dependent apoptosis. *Mol Neurobiol*, 56, 1-12, 2019. DOI: 10.1007/s12035-018-1083-z
3. Cheng W, Yang S, Liang F, Wang W, Zhou R, Li Y, Feng Y, Wang Y: Low-dose exposure to TCS disrupted osteogenic differentiation of mouse embryonic stem cells via BMP/ERK/Smad/Runx-2 signalling pathway. *Food Chem Toxicol*, 127, 1-10, 2019. DOI: 10.1016/j.fct.2019.02.038
4. Oliveira R, Domingues I, Koppe Grisolia C, Soares AMVM: Effects of triclosan on zebrafish early-life stages and adults. *Environ Sci Pollut Res*, 16, 679-688, 2009. DOI: 10.1007/s11356-009-0119-3
5. Weatherly LM, Gosse JA: Triclosan exposure, transformation, and

- human health effects. *J Toxicol Environ Health B Crit Rev*, 20 (8): 447-469, 2017. DOI: 10.1080/10937404.2017.1399306
- 6. Dhillon GS, Kaur S, Pulicharla R, Brar SK, Cleidon M, Verma M, Surampalli RY:** Triclosan: Current status, occurrence, environmental risks and bioaccumulation potential. *Int J Environ Res Public Health*, 12 (5): 5657-5684, 2015. DOI: 10.3390/ijerph120505657
- 7. Sandborgh-Englund G, Adolfsson-Erici M, Odham G, Ekstrand J:** Pharmacokinetics of Triclosan following oral ingestion in humans. *J Toxicol Environ Health A*, 69 (20): 1861-1873, 2006. DOI: 10.1080/15287390600631706
- 8. Allmyr M, Adolfsson-Erici M, McLachlan MS, Sandborgh-Englund G:** Triclosan in plasma and milk from Swedish nursing mothers and their exposure via personal care products. *Sci Total Environ*, 372, 87-93, 2006. DOI: 10.1016/j.scitotenv.2006.08.007
- 9. Fang JL, Stingley RL, Beland FA, Harrouk W, Lumpkins DL, Howard P:** Occurrence, efficacy, metabolism, and toxicity of Triclosan. *J Environ Sci Health C Environ Carcinog Ecotoxicol Rev*, 28, 147-171, 2010. DOI: 10.1080/10590501.2010.504978
- 10. Wu Y, Chitranshi P, Loukotková L, Gamboa da Costa G, Beland FA, Zhang J, Fang JL:** Cytochrome P450-mediated metabolism of Triclosan attenuates its cytotoxicity in hepatic cells. *Arch Toxicol*, 91 (6): 2405-2423, 2017. DOI: 10.1007/s00204-016-1893-6
- 11. Chaudhari U, Nemade H, Sureshkumar P, Vinken M, Ates G, Rogiers V, Hescheler J, Hengstler JG, Sachinidis A:** Functional cardiotoxicity assessment of cosmetic compounds using human-induced pluripotent stem cell-derived cardiomyocytes. *Arch Toxicol*, 92 (1): 371-381, 2018. DOI: 10.1007/s00204-017-2065-z
- 12. Park BK, Gonzales ELT, Yang SM, Bang M, Choi CS, Shin CY:** Effects of triclosan on neural stem cell viability and survival. *Biomol Ther (Seoul)*, 24 (1): 99-107, 2016. DOI: 10.4062/biomolther.2015.164
- 13. Chen X, Xu B, Han X, Mao Z, Chen M, Du G, Talbot P, Wang X, Xia Y:** The effects of Triclosan on pluripotency factors and development of mouse embryonic stem cells and zebrafish. *Arch Toxicol*, 89, 635-646, 2015. DOI: 10.1007/s00204-014-1270-2
- 14. Van Maele-Fabry G, Delhaise F, Picard JJ:** Morphogenesis and quantification of the development of post-implantation mouse embryos. *Toxicol In Vitro*, 4 (2): 149-156, 1990. DOI: 10.1016/0887-2333(90)90037-t
- 15. Augustine-Rauch K, Zhang CX, Panzica-Kelly JM:** *In vitro* developmental toxicology assays: A review of the state of the science of rodent and zebrafish whole embryo culture and embryonic stem cell assays. *Birth Defects Res C Embryo Today*, 90 (2): 87-98, 2010. DOI: 10.1002/bdrc.20175
- 16. Unur E, Ülger H, Ekinci N, Hacıaloğulları M, Ertekin T, Kılıç E:** Effect of anti-basic fibroblast growth factor (anti-bFGF) on *in vitro* embryonic development in rat. *Anat Histol Embryol*, 38, 241-245, 2009. DOI: 10.1111/j.1439-0264.2009.00927.x
- 17. Tekinarslan İ, Unur E, Ülger H, Ekinci N, Ertekin T, Hacıaloğulları M, Arslan S:** The effects of FGF-9 on *in vitro* embryonic development. *Balkan Med J*, 28, 18-22, 2011. DOI: 10.5174/tutfd.2009.02019.2
- 18. Ertekin T, Ülger H, Nisari M, Karaca Ö, Unur E, Şahin U, Elmalı F:** Effects of angiotensin on *in vitro* embryonic rat development. *Kafkas Univ Vet Fak Derg*, 17 (5): 843-847, 2011. DOI: 10.9775/kvfd.2011.4637
- 19. Nisari M, Ülger H, Unur E, Karaca O, Ertekin T:** Effect of interleukin 12 (IL-12) on embryonic development and yolk sac vascularisation. *Bratisl Lek Listy*, 115 (9): 532-537, 2014. DOI: 10.4149/blil_2014_103
- 20. Toder V, Carp H, Fein A, Torchinsky A:** The role of pro- and anti-apoptotic molecular interactions in embryonic maldevelopment. *Am J Reprod Immunol*, 48, 235-244, 2002. DOI: 10.1034/j.1600-0897.2002.01130.x
- 21. McArthur K, Kile BT:** Apoptotic caspases: Multiple or mistaken identities? *Trends Cell Biol*, 28 (6): 475-493, 2018. DOI: 10.1016/j.tcb.2018.02.003
- 22. Aşan E, Dağdeviren A:** Hücre. In, Aşan E, Dağdeviren A (Eds): *Moleküler Histoloji*. 145-168, Atlas Kitapçılık, Ankara, 2012.
- 23. New DAT:** Whole-embryo culture and the study of mammalian embryos during organogenesis. *Biol Rev Camb Philos Soc*, 53, 81-122, 1978.
- 24. Horie Y, Yamagishi T, Takahashi H, Iguchi T, Tatarazako N:** Effects of Triclosan on Japanese medaka (*Oryzias latipes*) during embryo development, early life stage and reproduction. *J Appl Toxicol*, 38 (4): 544-551, 2018. DOI: 10.1002/jat.3561
- 25. Ho JCH, Hsiao CD, Kawakami K, Tse WKF:** Triclosan (TCS) exposure impairs lipid metabolism in zebrafish embryos. *Aquat Toxicol*, 173, 29-35, 2016. DOI: 10.1016/j.aquatox.2016.01.001
- 26. Guo J, Ito S, Nguyen HT, Yamamoto K, Tanoue R, Kunisue T, Iwata H:** Effects of prenatal exposure to triclosan on the liver transcriptome in chicken embryos. *Toxicol Appl Pharmacol*, 347, 23-32, 2018. DOI: 10.1016/j.taap.2018.03.026
- 27. Haggard DE, Noyes PD, Waters KM, Tanguay RL:** Phenotypically anchored transcriptome profiling of developmental exposure to the antimicrobial agent, triclosan, reveals hepatotoxicity in embryonic zebrafish. *Toxicol Appl Pharmacol*, 308, 32-45, 2016. DOI: 10.1016/j.taap.2016.08.013
- 28. Szychowski KA, Sitarz AM, Wojtowicz AK:** Triclosan induces Fas receptor-dependent apoptosis in mouse neocortical neurons *in vitro*. *Neuroscience*, 284, 192-201, 2015. DOI: 10.1016/j.neuroscience.2014.10.001
- 29. Dubey D, Srivastav AK, Singh J, Chopra D, Qureshi S, Kushwaha HN, Singh N, Ray RS:** Photoexcited Triclosan induced DNA damage and oxidative stress via p38 MAP kinase signaling involving type I radicals under sunlight/UVB exposure. *Ecotoxicol Environ Saf*, 174, 270-282, 2019. DOI: 10.1016/j.ecoenv.2019.02.065
- 30. Lee GA, Choi KC, Hwang KA:** Treatment with phytoestrogens reversed triclosan and bisphenol A-induced anti-apoptosis in breast cancer cells. *Biomol Ther (Seoul)*, 26 (5): 503-511, 2018. DOI: 10.4062/biomolther.2017.160
- 31. O'Byrne KJ, Richard DJ:** Nucleolar caspase-2: Protecting us from DNA damage. *J Cell Biol*, 216 (6): 1521-1523, 2017. DOI: 10.1083/jcb.201704114
- 32. Ando K, Parsons MJ, Shah RB, Charendoff CI, Paris SL, Liu PH, Fassio SR, Rohrman BA, Thompson R, Oberst A, Sidi S, Bouchier-Hayes L:** NPM1 directs PIDDosome-dependent caspase-2 activation in the nucleolus. *J Cell Biol*, 216 (6): 1795-1810, 2017. DOI: 10.1083/jcb.201608095
- 33. Lamkanfi M, Kanneganti TD:** Caspase-7: A protease involved in apoptosis and inflammation. *Int J Biochem Cell Biol*, 42, 21-24, 2010. DOI: 10.1016/j.biocel.2009.09.013
- 34. Brentnall M, Rodriguez-Menocal L, De Guevara RL, Cepero E, Boise LH:** Caspase-9, caspase-3 and caspase-7 have distinct roles during intrinsic apoptosis. *BMC Cell Biol*, 14:32, 2013. DOI: 10.1186/1471-2121-14-32
- 35. Galluzzi L, López-Soto A, Kumar S, Kroemer G:** Caspases connect cell-death signaling to organismal homeostasis. *Immunity*, 44 (2): 221-231, 2016. DOI: 10.1016/j.immuni.2016.01.020
- 36. Li X, An J, Li H, Qiu X, Wei Y, Shang Y:** The methyl-triclosan induced caspase-dependent mitochondrial apoptosis in HepG2 cells mediated through oxidative stress. *Ecotoxicol Environ Saf*, 182:109391, 2019. DOI: 10.1016/j.ecoenv.2019.109391